



Tasmanian Digestive Centre

REFERRAL FOR LIVER FIBROSIS ULTRASOUND ASSESSMENT INCLUDING FIBROSCAN®

• Attach any current reports and investigations • Patient will be contacted with appointment details									
Enquiries:		Tasmanian Digestive Centre		T (03) 6449 4600					
Send referral to:		F 61399621090		E reception@tdctas.au					
Address:		75 - 79 Oldaker Street, Devonport, TAS, 7310							
Patient Details *mandatory fields									
Family Name*			Given Name*						
Date of Birth*			Sex ☐ Female ☐ Male ☐ Other						
Address*			Telephone*						
Medicare No			Reference No		Expiry				
☐ Bulk Bill ☐ DVA ☐ Other									
Interpreter		☐ Yes ☐ No		Language					
Aboriginal or Torres Strait Islander									
Cultural considerations / special needs									
Contact Person		Name							
		Relationship		Telephone					
Previous investigations									
Liver Biopsy		☐ Yes		Date		Fibrosis Stage ☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4		Inflammatory Grade ☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4	
Liver Function		☐ Yes		Date		Total Protein		Albumin	
						Bilirubin		ALT	
						GGT		ALP	
Haematology		☐ Yes		Date		Haemoglobin		Platelets	
FibroScan®		☐ Yes		Date		Result			
Indication		☐ Hepatitis B		☐ Hepatitis C		☐ NAFLD/MAFLD		☐ IDDM / NIDDM	
		☐ Alcohol		☐ PBC		☐ PSC		☐ Other <i>list</i>	
Clinician Estimate of Fibrosis		☐ No / minimal (FO-1)		☐ Moderate (F2-3)		☐ Severe / Cirrhosis (F4)			
Clinical Notes									
Referrer Details			Date of Referral			Provider No			
Name		Address							
Telephone		Fax							
Email		Copies to							

NOTE: This referral is valid for a FibroScan® assessment and an hepatic ultrasound which will be used to examine the right lobe of the liver, exclude vascular structures or other lesions which may interfere with the FibroScan® assessment and determine the skin-to-capsule distance to facilitate optimal FibroScan® probe selection. This comprehensive assessment is performed to optimize the liver fibrosis assessment and should not be considered to replace the patient's other abdominal imaging requirements or exclude other pathology. Patients will need to fast for four hours prior to undergoing FibroScan®. FibroScan® assessment may not be possible in up to 1/4 of patients with a BMI > 30kg / m² and alternative investigations may be more appropriate. For more information regarding the use of FibroScan or interpretation of results, contact Tasmanian Digestive Centre.